

Short explanation regarding filling in the order form / prescription

1. In order to ensure a responsible stop of medication use, a gradual tapering is often required. If tapering with the available registered dosages doesn't succeed, or results in too much withdrawal symptoms, tapering-medication (lower compounded dosages) can be prescribed.
2. Tapering can be done by two scientific proven methods:
 - * *Ashton Manual Curve* in which the tapering-traject is conform one entire hyperbolic traject (smaller steps towards the end), or
 - * *Hyperbolic reduction from registered dose to registered dose*. This type of tapering leads to small separate hyperbolic trajects from one dose to another.
3. **Cross box(es) risk factors** ; duration of medicine use; current dose; intake moment(s)
4. **Data of patient** needs to be filled in or attach label with name, address, city, country, phone number and email-address (mandatory due to communication with patient). Also mention name of local pharmacy.
5. **Data of prescriber (practitioner)** needs to be filled in or place stamp with the information.
6. Choose Tapering-traject in **Table 4**: select at least the first tapering-traject,
 - **or**, (if desired traject is not displayed in table) select the option **customized** and fill in the *start-* and *end dose*.
 - **or** (if patient needs to stabilize) select the option **Stabilizationstrip** and fill in the dose on which needs to be stabilized.
7. Mention date and have the practitioner sign and stamp the order form (= valid prescription). Then send the order form by fax or email to the number/email address mentioned on top of the page.
 - To request an advice regarding the tapering-traject, see **page 3**.
 - In case of a patient with a risk for withdrawal symptoms, go to **page 4** (Tapering with a 10% percentage per week) of 15% per week (=50% per month).
8. **Important** is the filling in of the self-monitoring form during the tapering-traject. This enables patients and practitioners to determine withdrawal symptoms a.s.a.p..
 In case of withdrawal symptoms: Aggravation of the withdrawal symptoms, or the failure of the tapering-traject can be prevented by having the patient stabilized (on the lowest dose with no withdrawal symptoms) and/or selecting a more gradual tapering-traject.

In order to realize a responsible tapering-traject, these strips can be prescribed:

Taperingstrip	Strip for 28 days to lower the daily dose; start- and end-dose are registered doses
Customized strip	Taperingstrip for 28 days with self chosen start- and end-dose (customized)
Stabilizationstrip	Strip for 28 days to stabilize a patient on the same dose (for 28 days)
Switchstrip	Strip for a responsible switch to another medicine in 28 days

Concepts

Taperingmedication is developed to provide an opportunity to taper/decrease the use of medication in a safe and responsible way. This traject can be customized and adjusted if necessary. The determination of the tapering-traject is done on the base of *shared decision making* and (self-)monitoring; this in accordance with accurate guidelines.

In a **Taperingstrip** the daily dose (intake) is decreased conform a hyperbolic curve. Hyperbolic means that the steps (difference in daily intake) toward the end become smaller. This is necessary in order to avoid/prevent the withdrawal symptoms as much as possible.

A **Taper-traject** can exist of one or more Taperingstrips, which are being used sequential; whether or not in combination with one or more stabilizationstrip(s).

A **Stabilizationstrip** can, when withdrawal symptoms occur, be prescribed in order to stabilize a patient on a certain dose. The patient can then stabilize on that dose before continuing the (adjusted) tapering-traject.

Selfmonitoring: attached with the tapering-medication a self-monitoring form is provided. This form needs to be filled in once a day so patient and practitioner can monitor the severity of the potential withdrawal symptoms (if they occur). If the severity increases, adjustments can be made to the tapering-traject in order to prevent withdrawal symptoms.

Instructions for the order form / prescription for Taperingstrips™

1. MEDICAL INDICATION: please mark all the indications applicable to the patient:
 - i. **preventive** reducing the risk of withdrawal symptoms or relapse effects caused by diminishing the dose too quickly.
 - ii. **reduction anxiety** patient is excessively anxious about tapering, for example as a result of comorbid anxiety disorder.
 - iii. **failed attempt** previous attempts to stop or reduce the dose have failed.
 - iv. **distinction** distinguishing between withdrawal symptoms and possible relapse.
 - v. **slow metabolizer** the lowest dose causes a high plasma concentration, increasing the risk of withdrawal effects.
 - vi. **high dose** the patient has used the medication for a period of 6 months at a dose at least 150% of the defined daily dose.
2. Fill in the patient's details.
3. Fill in the details of the treating physician.
4. Fill in the table. Below you can see four examples showing the different ways of filling in the table:

a. **Total reduction**, using **3** tapering strips™: mark **3** boxes (schedule 3 x 28 days = 84 days)

starting dose EXAMPLE	taper duration	STRIP 1 28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days
30 mg/day	112 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-5	<input type="checkbox"/> EXMPL 5-2	<input type="checkbox"/> EXMPL 2-0
	84 days	<input checked="" type="checkbox"/> EXMPL 30-10	<input checked="" type="checkbox"/> EXMPL 10-5	<input checked="" type="checkbox"/> EXMPL 5-0	
	56 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-0		

b. **Dose reduction** using **2** tapering strips™: mark **2** boxes (schedule 2 x 28 days = 56 days)

starting dose EXAMPLE	taper duration	STRIP 1 28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days
30 mg/day	112 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-5	<input type="checkbox"/> EXMPL 5-2	<input type="checkbox"/> EXMPL 2-0
	84 days	<input checked="" type="checkbox"/> EXMPL 30-10	<input checked="" type="checkbox"/> EXMPL 10-5	<input type="checkbox"/> EXMPL 5-0	
	56 days	<input type="checkbox"/> EXMPL 30-10	<input type="checkbox"/> EXMPL 10-0		

c. **Customized schedule** please mark each box, and fill in the desired start and end dose for each strip. Total duration = number of strips x 28 days.

CUSTOMIZED 2 x 28 days	<input checked="" type="checkbox"/> EXMPL from 60 - 30 mg	<input checked="" type="checkbox"/> EXMPL from 30 - 10 mg
-------------------------------	---	---

d. **Stabilization** to a steady dose: mark the box, and fill in daily dose and number of repeats (e.g.: 17 mg/day and 2x repeats=> schedule (1 + 2) x 28 days = 84 days)

STABILIZATION 28 days	<input checked="" type="checkbox"/> EXMPL stab.	17 mg/day	2 repeats
------------------------------	---	------------------	------------------

5. Sign the prescription.
6. Fax the prescription to the Regenboog Apotheek: **+31-(0)85-2736129** or scan and mail to tapering@regenboogapotheek.com

NOTE: Please only fax the order form / prescription

Do not fax the instruction sheets.

Request for advice QUETIAPINE 

Regenboog Apotheek has guided more than 10.000 patients before and during the tapering-traject. Advice regarding the traject will be send within three working days.

Dear sir / madam,

I, arts / patient (cross box), would like to receive an advice for a tapering-traject for the patiënt mentioned underneath. Patient has the following characteristics:

- * Patient received quetiapine for the indication of:
- * Patient is currently free of symptoms: yes no
- * The reason(s) for tapering is/are:

See explanation page 6, cross all boxes which are applicable.

1a. Risk factor: missed intake fear tapering previous failed attempts distinguish slow metaboliser high dose start others

1b. Duration of use of quetiapine: <1 year 1-2 year 2-5 year 5-10 year >10 year

1c. Current intake quetiapine:, mg time hour decrease desired end dose.....
, mg time hour decrease desired end dose.....
, mg time hour decrease desired end dose.....
, mg time hour decrease desired end dose.....

1d. Other oral medication: (name), mg per day
 (name), mg per day
 (name), mg per day
 (name), mg per day
 (name), mg per day

1e. Tablets are the desired form of medication (cross box)

1f. Required tapering period: .. months (number) **or**
 10% per week (=30% per month)
 15% per week (=50% per month)

2. Initials and surname patient:

Date of birth (dd-mm-yyyy): .. - .. - .. Gender: M / F

Street and number:

Postal code and city: Country:

E-mail address (**mandatory**): Phone:

3. Name practitioner:

Medical registration number:

Street and number of practice:

Postal code and city:

E-mail address (**mandatory**):

Phone:

Date: Stamp practitioner (preferably):

Signature practitioner and/or patient:
 I hereby declare that all information has been provided truthfully

Shipping address: Regenboog Apotheek, Antwoordnummer 16500, 4840 WJ BAVEL, Nederland

Order form and prescription

R/ Order form = Prescription for QUETIAPINE Taperingstrips™ ATC code: N05AH04

1a. Medical indication: preventive reduction anxiety failed attempt distinction slow metabolizer high dose

1b. Duration of quetiapine use: <1 year 1-2 years 2-5 years 5-10 years >10 years

2. Full name of patient:

Date of birth (dd-mm-yyyy): . . . - . . . - gender: M / F

Address:

City, postal code and country:

E-mail address (**mandatory**): Phone:

3. Name of treating physician:

Medical registration number:

Address:

Postal code and country:

E-mail address (**mandatory**):

Phone:

4. Please, fill in the **table A** (Asthon Manual schedule), following the instructions above, **use the startdose (mg) of the intake dose !**

Startdose quetiapine	Tapering period	QTPN=quetiapine STRIP 1=28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days	STRIP 5 28 days
800 mg/Intake10%	280 days	<input type="checkbox"/> QTPN 800-535	<input type="checkbox"/> QTPN 535-350	<input type="checkbox"/> QTPN 535-225	<input type="checkbox"/> QTPN 225-150	<input type="checkbox"/> QTPN 150-100→
800 mg/Intake15%	196 days	<input type="checkbox"/> QTPN 800-400	<input type="checkbox"/> QTPN 400-200	<input type="checkbox"/> QTPN 200-100	<input type="checkbox"/> QTPN 100-50→	
600 mg/Intake10%	280 days	<input type="checkbox"/> QTPN 600-400	<input type="checkbox"/> QTPN 400-265	<input type="checkbox"/> QTPN 265-175	<input type="checkbox"/> QTPN 175-120	<input type="checkbox"/> QTPN 120-80→
600 mg/Intake15%	196 days	<input type="checkbox"/> QTPN 600-300	<input type="checkbox"/> QTPN 300-150	<input type="checkbox"/> QTPN 150-80→		
400 mg/Intake10%	252 days	<input type="checkbox"/> QTPN 400-265	<input type="checkbox"/> QTPN 265-175	<input type="checkbox"/> QTPN 175-120	<input type="checkbox"/> QTPN 120-80→	
400 mg/Intake15%	168 days	<input type="checkbox"/> QTPN 400-200	<input type="checkbox"/> QTPN 200-100	<input type="checkbox"/> QTPN 100-50→		
300 mg/Intake10%	224 days	<input type="checkbox"/> QTPN 300-200	<input type="checkbox"/> QTPN 200-145	<input type="checkbox"/> QTPN 145-100→		
300 mg/Intake15%	140 days	<input type="checkbox"/> QTPN 300-150	<input type="checkbox"/> QTPN 150-75	<input type="checkbox"/> QTPN 75-30	<input type="checkbox"/> QTPN 30-15	<input type="checkbox"/> QTPN 15-0
100 mg/Intake10%	140 days	<input type="checkbox"/> QTPN →100-60	<input type="checkbox"/> QTPN 60-40	<input type="checkbox"/> QTPN 40-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0
100 mg/Intake15%	112 days	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0	
80 mg/Intake10%	140 days	<input type="checkbox"/> QTPN →80-55	<input type="checkbox"/> QTPN 55-35	<input type="checkbox"/> QTPN 35-20	<input type="checkbox"/> QTPN 20-10	<input type="checkbox"/> QTPN 10-0
80 mg/Intake15%	112 days	<input type="checkbox"/> QTPN →80-40	<input type="checkbox"/> QTPN 40-20	<input type="checkbox"/> QTPN 20-10	<input type="checkbox"/> QTPN 10-0	
50 mg/Intake10%	112 days	<input type="checkbox"/> QTPN →50-35	<input type="checkbox"/> QTPN 35-20	<input type="checkbox"/> QTPN 20-10	<input type="checkbox"/> QTPN 10-0	
50 mg/Intake15%	84 days	<input type="checkbox"/> QTPN →50-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0		
customized	. X 28 days	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN
rescue	3 X 15 tablets	<input type="checkbox"/> QTPN 1 mg				
stable	28 days	<input type="checkbox"/> QTPN stable	. . . mg	.. repeats	dose round off to 1 mg	<input type="checkbox"/> send immediately

COSTS Tapering- and stabilization strips contain medication for a fixed period of 28 days, all strips have the same price. Taperingstrip™: € 77.-; Stabilization strip € 38.50 for a period of 28 days; Prices are subject to change.

5. Date: Stamp of treating physician

Signature of treating physician:

Shipping address: **Regenboog Apotheek, Antwoordnummer 16500, 4840 WJ BAVEL, The Netherlands**

After submitting the form, fax to Regenboog Apotheek +31-(0)85-2736129
or scan and mail to tapering@regenboogapothek.com

Order form and prescription

R/ Order form = prescription for QUETIAPINE Taperingstrips™ ATC code: N05AH04

1a. Medical indication: preventive reduction anxiety failed attempt distinction slow metabolizer high dose

1b. Duration of quetiapine use: <1 year 1-2 years 2-5 years 5-10 years >10 years

2. Full name of patient:

Date of birth (dd-mm-yyyy): . . . - . . . - gender: M / F

Address:

City, postal code and country:

E-mail address (mandatory): Telephone no:

3. Name of treating physician:

Medical registration number:

Address:

Postal code and country:

E-mail address (mandatory):

Telephone no:

4. Please, fill in the **table B** (Regular schedule), following the instructions above, **use the startdose (mg) of the in-takedose !** **Most prescribed Trajectories**

Startdose quetiapine	Tapering period	QTPN=quetiapine STRIP 1=28 days	STRIP 2 28 days	STRIP 3 28 days	STRIP 4 28 days	STRIP 5 28 days
300 mg/Intake	140 days	<input type="checkbox"/> QTPN 300-200	<input type="checkbox"/> QTPN 200-100	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-0
	140 days	<input type="checkbox"/> QTPN 300-100	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0
200 mg/Intake	112 days	<input type="checkbox"/> QTPN 200-100	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0
	112 days	<input type="checkbox"/> QTPN 200-100	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-0	
100 mg/Intake	84 days	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0	
	84 days	<input type="checkbox"/> QTPN 100-50	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-0		
	56 days	<input type="checkbox"/> QTPN 100-25	<input type="checkbox"/> QTPN 25-0			
50 mg/Intake	84 days	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0		
	56 days	<input type="checkbox"/> QTPN 50-25	<input type="checkbox"/> QTPN 25-0			
25 mg/Intake	84 days	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-5	<input type="checkbox"/> QTPN 5-0		
	56 days	<input type="checkbox"/> QTPN 25-10	<input type="checkbox"/> QTPN 10-0			
	28 days	<input type="checkbox"/> QTPN 25-0				
customized	. X 28 days	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN	<input type="checkbox"/> QTPN
set-up	1 X . . days	<input type="checkbox"/> QTPN				
rescue	3X 15 tablets	<input type="checkbox"/> QTPN 1 mg				
stable	28 days	<input type="checkbox"/> QTPN stable	. . . mg	.. repeats	dose round off to 1 mg	<input type="checkbox"/> send immediately

COSTS Tapering- and stabilization strips contain medication for a fixed period of 28 days, all strips have the same price. Taperingstrip™: € 77.-; Stabilization strip € 38.50 for a period of 28 days; Prices are subject to change.

5. Date: Stamp of treating physician

Signature of treating physician:

Shipping address: **Regenboog Apotheek, Antwoordnummer 16500, 4840 WJ BAVEL, The Netherlands**

Extra Instruction for QUETIAPINE (Seroquel)

Please ensure you have the correct prescription and order form in front of you. There are several different order-forms for Antidepressants (38), Anti-psychotics (39), Sedatives (27), Anti-epileptics (13), Hormones (2) & Analgetics (4).

PLEASE READ CAREFULLY BEFORE FILLING IN THE ORDER FORM / PRESCRIPTION

1. A sudden termination of treatment with quetiapine, or a sudden reduction in dosage can lead to significant withdrawal symptoms. To avoid this, the intake dose should be reduced in small steps gradually over a period of time. This can be done using Taperingstrips™.
2. One or more Taperingstrips™ can be used to gradually reduce the dose of quetiapine to a desired lower dose or to zero. If more than one strip is needed the final dose of the previous strip is always equal to the starting dose of the following strip. This has to be done for every intake moment(if there are more than one intakes a day).
3. The treating physician (psychiatrist or general practitioner) should plan the tapering schedule together with the patient. It is not possible to predict the perfect schedule for each patient. Some patients are more sensitive to withdrawal symptoms than others, and some patients cope with symptoms better than others. The risk of withdrawal symptoms and their intensity can be reduced by choosing a more gradual tapering schedule. There are two options when choosing a tapering schedule:
 - 1) **Choose a tapering schedule according the Asthon Manual for a fixed percentage (10 or 15% a week).**
 - a. **Variable** - Choose a tapering schedule for the first 28 days only, then use the patient's experiences and feedback (from the first three weeks) to inform the physician about the remainder of the tapering trajectory. To do this only the **black** part of **table A** on page 4 has to be filled in. The experiences of the patient during the first weeks (withdrawal symptoms or not) can then be used to choose the tapering trajectories for the following strip(s) by filling in in the **red** part of the **table A** on page 4 you need several tapering trajectories (→).
 - b. **Fixed** - Choose a tapering schedule for the entire period of the dosage reduction. To do this, fill in both the **black** and (if applicable) the **red** part of **table A** on page 5 you need several tapering trajectories (→).
 - 2) **Choose a tapering schedule from (start) and to (end) doses which are on the market (Regular Schedule)**
 - a. **Variable** - Choose a tapering schedule for the first 28 days only, then use the patient's experiences and feedback (from the first three weeks) to inform the physician about the remainder of the tapering trajectory. To do this only the **black** part of **table B** on page 5 has to be filled in. The experiences of the patient during the first weeks (withdrawal symptoms or not) can then be used to choose the tapering trajectories for the following strip(s) by filling in in the **red** part of the **table B** on page 5.
 - b. **Fixed** - Choose a tapering schedule for the entire period of the dosage reduction. To do this, fill in both the **black** and (if applicable) the **red** part of **table B** on page 5.
4. Quetiapine Taperingstrips™ are only available on prescription. The (by physician) signed order-form is seen as a valid prescription. After filling in the form, the treating physician should terminate any current prescription for quetiapine at the local pharmacy.
5. In most countries, the costs of the medication in the Taperingstrips™ are covered by health insurance. Tapering- and stabilizationstrips contain medication for a fixed period of 28 days, all strips have the same price. Taperingstrip™: € 77.-; Stabilization-strip € 38.50 for a period of 28 days. Prices are subject to change, so please check before ordering.
6. Once the order-form/prescription has been received by the Regenboog Apotheek, we will send an invoice. Once we've received the payment, the Taperingstrip™ will be shipped to the patient's home address. The patient will receive an invoice and explanation beforehand by mail.
7. A Taperingstrip™ contains 28 numbered pouches, with each pouch containing one or several tablets. The number of pills may vary per day. The total dose of the medication reduces over time as the patient takes the pills in each daily pouch.
8. The patient starts their tapering schedule (preferably) on a Sunday by taking the pills in the first pouch, numbered 028. **By counting down**, the patient finishes four weeks later on a Saturday, by taking the pills in the last pouch, numbered 001. By doing this, the patient can easily keep track of how many doses and days are left.
9. Each pouch of the strip is printed with the sequence number, the colour and strength of each tablet, the total number of tablets and the name of the pharmacy.
10. **STABILIZATION:** A stabilizationstrip can keep the patient on a fixed dose for a desired period of time. Such stabilization may be necessary if tapering appears to be going too fast for the patient, or the patient does not wish to reduce the dose to zero, but remain at an optimal lower dose. The stabilization-strip can be produced at almost any desired dose rounded off to 1 mg quetiapine.
11. In some cases, the patient is currently using a dose that is higher or different from the starting dose of the standard available Taperingstrips (see table 4 on page 4). These patients can also reduce their dose gradually by using **customized** Taperingstrips™ for the first part of the taper (see table 4 on page 4).

12. For the **Asthon Manual** schedule quetiapine, the following Taperingstrips™ are standard available:

Startdose quetiapine	Enddose quetiapine	Tapering period	Name of the strip QTPN=quetiapine	Step dose reduction	average dose reduction	average dose reduction
600mg/dose	400mg/dose	28 days	QTPN 600-400/28days	1 mg	7,14mg/dose	10%/week
600mg/dose	300mg/dose	28 days	QTPN 600-300/28days	1 mg	10,71mg/dose	15%/week
300mg/dose	200mg/dose	28 days	QTPN 300-200/28days	1 mg	3,57mg/dose	10%/week
300mg/dose	150mg/dose	28 days	QTPN 300-150/28days	1 mg	5,36mg/dose	15%/week
150mg/dose	100mg/dose	28 days	QTPN 150-100/28days	1 mg	1,79mg/dose	10%/week
150mg/dose	80mg/dose	28 days	QTPN 150-80/28days	1 mg	2,5mg/dose	14%/week
100mg/dose	60mg/dose	28 days	QTPN 100-60/28days	1 mg	1,43mg/dose	12%/week
80mg/dose	40mg/dose	28 days	QTPN 80-40/28days	1 mg	1,43mg/dose	15%/week
60mg/dose	40mg/dose	28 days	QTPN 60-40/28days	1 mg	0,71mg/dose	10%/week
50mg/dose	35mg/dose	28 days	QTPN 50-35/28days	1 mg	0,54mg/dose	9%/week
50mg/dose	25mg/dose	28 days	QTPN 50-25/28days	1 mg	0,89mg/dose	15%/week
40mg/dose	25mg/dose	28 days	QTPN 40-25/28days	1 mg	0,54mg/dose	12%/week
40mg/dose	20mg/dose	28 days	QTPN 40-20/28days	1 mg	0,71mg/dose	15%/week
35mg/dose	20mg/dose	28 days	QTPN 35-20/28days	1 mg	0,54mg/dose	13%/week
25mg/dose	10mg/dose	28 days	QTPN 25-10/28days	1 mg	0,54mg/dose	18%/week
20mg/dose	10mg/dose	28 days	QTPN 20-10/28days	1 mg	0,36mg/dose	15%/week
10mg/dose	zero	28 days	QTPN 10-0/28days	1 mg	0,36mg/dose	30%/week
stable	. . .mg/dose	28 days	QTPN stable	0 mg	no reduction	0%/week

The table below shows how many mg of quetiapine each daily pouch contains in the **Asthon Manual** schedule.

Schedule: quetiapine		600 - 400	600 - 300	300 - 200	300 - 150	150 - 100	150 - 80	100 - 60	80 - 40	60 - 40	50 - 35	50 - 25	40 - 25	40 - 20	35 - 20	25 - 10	20 - 10	10 - 0
Sachet No	Intake day	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg	mg
sachet 28-01	Sunday	600	600	300	300	150	150	100	80	60	50	50	40	40	35	25	20	10
sachet 27-01	Monday	590	581	295	291	148	146	98	78	59	49	49	39	39	34	24	20	10
sachet 26-01	Tuesday	580	563	290	283	145	143	96	76	58	49	48	39	38	34	24	19	9
sachet 25-01	Wednesday	570	546	286	275	143	140	95	74	57	48	47	38	37	33	23	19	9
sachet 24-01	Thursday	561	530	281	267	141	136	93	73	57	48	46	38	37	33	23	18	8
sachet 23-01	Friday	552	515	277	260	139	133	91	71	56	47	45	37	36	32	22	18	8
sachet 22-01	Saturday	543	500	272	253	137	130	89	69	55	46	44	36	35	31	21	18	8
sachet 21-01	Sunday	534	486	268	246	135	127	88	68	54	46	43	36	34	31	21	17	7
sachet 20-01	Monday	526	473	264	240	133	124	86	66	54	45	42	35	33	30	20	17	7
sachet 19-01	Tuesday	518	460	260	233	131	121	85	64	53	45	41	35	33	30	20	16	6
sachet 18-01	Wednesday	510	448	256	227	129	119	83	63	52	44	40	34	32	29	19	16	6
sachet 17-01	Thursday	502	436	252	221	127	116	82	61	51	44	39	33	31	28	18	16	6
sachet 16-01	Friday	494	425	248	216	125	113	80	60	50	43	38	33	30	28	18	15	5
sachet 15-01	Saturday	487	414	245	210	123	111	79	58	50	42	37	32	30	27	17	15	5
sachet 14-01	Sunday	480	404	241	205	121	108	77	57	49	42	36	32	29	27	17	15	5
sachet 13-01	Monday	473	394	237	200	120	106	76	55	48	41	35	31	28	26	16	14	4
sachet 12-01	Tuesday	466	385	234	195	118	103	74	54	48	41	34	31	27	26	16	14	4
sachet 11-01	Wednesday	459	375	231	190	116	101	73	53	47	40	33	30	27	25	15	14	3
sachet 10-01	Thursday	453	367	227	186	114	99	71	51	46	40	32	30	26	25	15	13	3
sachet 09-01	Friday	446	358	224	181	113	96	70	50	45	39	31	29	25	24	14	13	3
sachet 08-01	Saturday	440	350	221	177	111	94	69	49	45	39	31	29	25	24	14	12	2
sachet 07-01	Sunday	434	342	218	173	109	92	67	47	44	38	30	28	24	23	13	12	2
sachet 06-01	Monday	428	334	215	169	108	90	66	46	43	38	29	28	23	23	13	12	2
sachet 05-01	Tuesday	422	327	212	165	106	88	65	45	43	37	28	27	23	22	12	11	1
sachet 04-01	Wednesday	416	320	209	161	105	86	64	44	42	37	27	27	22	22	11	11	1
sachet 03-01	Thursday	411	313	206	157	103	84	62	42	41	36	27	26	21	21	11	11	1
sachet 02-01	Friday	405	306	203	154	101	82	61	41	41	36	26	25	21	20	10	10	1
sachet 01-01	Saturday	400	300	200	150	100	80	60	40	40	35	25	25	20	20	10	10	1

13. For the **Regular Schedule** quetiapine, the following Taperingstrips™ are standard available:

Startdose quetiapine	Enddose quetiapine	Tapering period	Name of the strip QTPN=quetiapine	Step dose reduction	average dose reduction	average dose reduction
300mg/dose	200mg/dose	28 days	QTPN 300-200/28days	4/3 mg	3,57mg/dose	10%/week
200mg/dose	100mg/dose	28 days	QTPN 200-100/28days	4/3 mg	3,57mg/dose	15%/week
100mg/dose	50mg/dose	28 days	QTPN 100-50/28days	2/1 mg	1,79mg/dose	15%/week
100mg/dose	25mg/dose	28 days	QTPN 100-25/28days	3/2 mg	2,68mg/dose	23%/week
50mg/dose	25mg/dose	28 days	QTPN 50-25/28days	1 mg	0,89mg/dose	15%/week
25mg/dose	10mg/dose	28 days	QTPN 25-10/28days	1 mg	0,54mg/dose	18%/week
25mg/dose	zero	28 days	QTPN 25-0/28days	1 mg	0,89mg/dose	30%/week
10mg/dose	zero	28 days	QTPN 10-0/28days	1 mg	0,36mg/dose	30%/week
stable	...mg/dose	28 days	QTPN stable	0 mg	no reduction	0%/week

The table below shows how many mg of quetiapine each daily pouch contains in the **Regular Schedule**.

Schedule: quetiapine		300 - 200	200 - 100	100 - 50	100 - 25	50 - 25	25 - 15	25 - 10	25 - 0	10 - 0
Sachet No	Intake day	mg	mg	mg	mg	mg	mg	mg	mg	mg
sachet 28-01	Sunday	300	200	100	100	50	25	25	25	10
sachet 27-01	Monday	295	195	98	96	49	25	24	24	10
sachet 26-01	Tuesday	290	189	95	91	48	24	24	23	9
sachet 25-01	Wednesday	286	184	93	87	47	24	23	22	9
sachet 24-01	Thursday	281	179	91	84	46	23	23	21	8
sachet 23-01	Friday	277	175	88	80	45	23	22	19	8
sachet 22-01	Saturday	272	170	86	76	44	23	21	18	8
sachet 21-01	Sunday	268	166	84	73	43	22	21	17	7
sachet 20-01	Monday	264	161	82	70	42	22	20	16	7
sachet 19-01	Tuesday	260	157	80	67	41	21	20	15	6
sachet 18-01	Wednesday	256	153	78	64	40	21	19	14	6
sachet 17-01	Thursday	252	149	76	61	39	21	18	13	6
sachet 16-01	Friday	248	145	74	58	38	20	18	13	5
sachet 15-01	Saturday	245	142	72	55	37	20	17	12	5
sachet 14-01	Sunday	241	138	71	53	36	20	17	11	5
sachet 13-01	Monday	237	135	69	50	35	19	16	10	4
sachet 12-01	Tuesday	234	131	67	48	34	19	16	9	4
sachet 11-01	Wednesday	231	128	65	45	33	19	15	8	3
sachet 10-01	Thursday	227	125	64	43	32	18	15	7	3
sachet 09-01	Friday	224	122	62	41	31	18	14	6	3
sachet 08-01	Saturday	221	119	60	38	31	17	14	5	2
sachet 07-01	Sunday	218	116	59	36	30	17	13	5	2
sachet 06-01	Monday	215	113	57	34	29	17	13	4	2
sachet 05-01	Tuesday	212	110	56	32	28	16	12	3	1
sachet 04-01	Wednesday	209	108	54	30	27	16	11	2	1
sachet 03-01	Thursday	206	105	53	29	27	16	11	2	1
sachet 02-01	Friday	203	103	51	27	26	15	10	1	1
sachet 01-01	Saturday	200	100	50	25	25	15	10	1	1


IMPORTANT NOTICE: Taperingstrips™ are not intended to replace medical care provided by a physician. Careful counselling and monitoring of the patient remains essential during and after treatment with tapering strips.

Please note: Treatment with the Taperingstrip™ supports the tapering of medication, reducing withdrawal effects to a minimum. If withdrawal effects do occur, they will be not as intensive or prolonged as during a traditional tapering approach. This will make it easier for the treating physician to recognize a relapse (such as a return of depression or anxiety) and to distinguish between relapse or withdrawal effects in good time.

There is a noticeable difference in timing between the occurrence of withdrawal symptoms and relapse. Withdrawal symptoms usually occur shortly or immediately after the start of tapering. Relapse usually occurs at a later stage, often after tapering with quetiapine has been completed. While withdrawal symptoms become less severe and disappear over time, symptoms as a result of relapse tend to remain and may even get worse.

Should you have any questions or if you require more information, please Regenboog Apotheek (Rainbow Pharmacy) at tapering@regenboogapothek.com or + 31-(0)161-437 137.

14. Tabel Effect* as a function of the daily change (decrease) in dose from 25 mg till 0 mg of the medicine in 28 days.



* "Effect" show the occupancy rate of the receptor (in percentages) where the medicine attaches as a function of the daily dose; from the lowest registered dose till 0. Change (decrease) should be as **gradual** as possible in order to prevent withdrawal symptoms as much as possible. This is will be realized by lowering the dose on a conform a hyperbolic function. Hyperbolic means that the steps (change in daily intake) toward the end will get smaller.

15. Important: Tapering medication Taperingstrips™ does not replace the usual care/consult of the practitioner! Good guidance of the patient during, but also after tapering remains essential in order to recognize a potential relapse.

16. Selfmonitoring-form in order to realize a responsible tapering-traject.

Attached with the tapering-medication a self-monitoring form is provided. This form needs to be filled in once a day so patient and practitioner can monitor the severity of the potential withdrawal symptoms (if they occur). If the severity increases, adjustments can be made to the tapering-traject in order to prevent withdrawal symptoms. Filling in the form does only take a minute per day.

17. Research to the responsible tapering of medication

Patients, who give there permission, can send the filled in self-monitoring form back to the pharmacy (free of costs). All data will be anonymized and prof. Jim van Os and dr. Peter C. Groot (University of Utrecht) will use the data for further research.



I.M.W. de Jong, J.C. van Vliet, apothekers
Brigidastraat 12, 4854 CT Bavel

Tel: (0161) 437 137
Fax: (0161) 437 138

KWITANTIE VOOR: DATUM: 12-11-2019

E Example
Brigidastraat 12
4854 CT Bavel

Code/Arts	Zi-nr	Omschrijving	Materiaal	Tarief	WMG	Bijbetaling	Bedrag	BTW
Betreft E Example, 31-12-1960/V								
S /SPBUI	1	ST PAROXETINE 5-OMG TAPERING		77,00			77,00	

Voorschrift van:

H = Huisarts
S = Specialist
V = Verloskundige
T = Tandarts

HS = Huisarts op advies specialist
HV = Handverkoop
P = Psychiater / Neuroloog

Bank NL70RABO039.50.37.697
KvK 20060757
BTW NL0087.63.914.B.01

KRB01792

RECEPTNUMMER	TE BETALEN
3722099P/KRB01792	77,00

Algemene Verkoop- en Betalingsvoorwaarden zijn op 1 februari 2009 gedeponereerd bij de Kamer van Koophandel Haaglanden te 's-Gravenhage (HR 40 409 373)



I.M.W. de Jong, J.C. van Vliet, apothekers
Brigidastraat 12, 4854 CT Bavel

Tel: (0161) 437 137
Fax: (0161) 437 138

KWITANTIE VOOR: DATUM: 12-11-2019

E Example
Brigidastraat 12
4854 CT Bavel

Code/Arts	Zi-nr	Omschrijving	Materiaal	Tarief	WMG	Bijbetaling	Bedrag	BTW
Betreft E Example, 31-12-1960/V								
S /SPBUI	1	ST PAROXETINE 5-OMG TAPERING		77,00			77,00	

DUPLIKAAT

KRB01792

RECEPTNUMMER	TE BETALEN
3722099P/KRB01792	77,00

Voorschrift van:

H = Huisarts
S = Specialist
V = Verloskundige
T = Tandarts

HS = Huisarts op advies specialist
HV = Handverkoop
P = Psychiater / Neuroloog

Bank NL70RABO039.50.37.697
KvK 20060757
BTW NL0087.63.914.B.01

Algemene Verkoop- en Betalingsvoorwaarden zijn op 1 februari 2009 gedeponereerd bij de Kamer van Koophandel Haaglanden te 's-Gravenhage (HR 40 409 373)

Invoice specification

Taperingstrip™ Paroxetine 5-0/28 contains the following medication:

invoice specification/Rechnungsangabe/spécification de la facture			
Number of tablets	Name medication	Dose	ATC Code
1	paroxetine	5,0 mg	N06AB05
14	paroxetine	2,0 mg	N06AB05
10	paroxetine	1,0 mg	N06AB05
12	paroxetine	0,50 mg	N06AB05
22	paroxetine	0,20 mg	N06AB05
12	paroxetine	0,10 mg	N06AB05

NR zakje	Schedule	mg totaal	Total number of tablets	5 mg	2 mg	1 mg	0,5 mg	0,2 mg	0,1 mg
	day / Tag / jour								
28	sunday, Sonntag, dimanche	5	1	1					
27	monday, Montag, lundi	4,6	4		2		1		1
26	tuesday, Dienstag, mardi	4,3	4		2			1	1
25	wednesday, Mittwoch, mercredi	4	2		2				
24	thursday, Donnerstag, jeudi	3,7	4		1	1	1	1	
23	friday, Freitag, vendredi	3,4	4		1	1		2	
22	saturday, Samstag, samedi	3,2	3		1	1		1	
21	sunday, Sonntag, dimanche	2,9	4		1		1	2	
20	monday, Montag, lundi	2,7	3		1		1	1	
19	tuesday, Dienstag, mardi	2,5	2		1		1		
18	wednesday, Mittwoch, mercredi	2,3	3		1			1	1
17	thursday, Donnerstag, jeudi	2,1	2		1				1
16	friday, Freitag, vendredi	1,9	4			1	1	2	
15	saturday, Samstag, samedi	1,8	4			1	1	1	1
14	sunday, Sonntag, dimanche	1,6	3			1	1		1
13	monday, Montag, lundi	1,4	3			1		2	
12	tuesday, Dienstag, mardi	1,3	3			1		1	1
11	wednesday, Mittwoch, mercredi	1,1	2			1			1
10	thursday, Donnerstag, jeudi	1	1			1			
9	friday, Freitag, vendredi	0,9	3				1	2	
8	saturday, Samstag, samedi	0,7	2				1	1	
7	sunday, Sonntag, dimanche	0,6	2				1		1
6	monday, Montag, lundi	0,5	1				1		
5	tuesday, Dienstag, mardi	0,4	2					2	
4	wednesday, Mittwoch, mercredi	0,3	2					1	1
3	thursday, Donnerstag, jeudi	0,2	1					1	
2	friday, Freitag, vendredi	0,1	1						1
1	saturday, Samstag, samedi	0,1	1						1
			71	1	14	10	12	22	12

(NL) Vergoeding conform C-120/95 en C-158/96 van het Hof van Justitie van de Europese Gemeenschap (Decker/Kohll-arrest)

De door de arts voorgeschreven farmacotherapeutische behandeling is overeenkomstig het behandelprotocol voor het afbouwen van paroxetine (Seroxat).

De patiënt heeft van de Regenboog Apotheek paroxetine (Seroxat) afbouwmedicatie ontvangen en heeft zijn/haar medicatie geleidelijk afgebouwd volgens schema.

De kosten van de medicatie in Nederland (€ 77,-) staan vermeld op de factuur.

Volgens het Decker/Kohll-arrest (C-120/95 en C-158/06) heeft de patiënt recht op vergoeding van de medicatie volgens het in zijn thuisland geldende tarief.

(GB) Repayment in accordance with C-120/95 C-158/96 (Decker Kohll judgment) of the Court of Justice of the European Union

The pharmaco therapeutic treatment prescribed by the doctor is in compliance with the treatment protocol for medication reduction paroxetine (Seroxat).

The patient received this treatment for medication reduction paroxetine (Seroxat) from pharmacy Regenboog Apotheek and he/she reduced his/her medication according to the schedule.

The cost of the medication in the Netherlands (€ 77,-) is listed on the invoice.

According to the Decker Kohll judgment (C-120/95 and C-158/06), the patient may claim repayment of the medication at the rate applicable to his homeland.

(DE) Erstattung nach C-120/95 und C-158/96 (Urteil Decker und Kohll) des Gerichtshofs der Europäischen Union

Diese vom Arzt verschrieben pharmatherapeutische Behandlung stimmt überein mit dem Behandlungsprotokoll für Medizinabbau paroxetine (Seroxat).

Der Patient/die Patientin erhielt diese Abbaumedikamenten paroxetine (Seroxat) vom Regenboog Apotheek und hat seine/ihre Medikamente reduziert nach dem Zeitplan.

Die Kosten für die Medikamente in den Niederlanden (€ 77,-) sind auf der Rechnung aufgeführt.

Nach dem Urteil (C-120/95 und C-158/06, Decker und Kohll) hat der Patient Anspruch auf Erstattung der Medikamente nach dem Kurs in seiner Heimat.

(FR) Remboursement conform C-120/95 et C-158/96 (l'arrêt Decker Kohll) de la Cour de justice de l'Union européenne

Le traitement pharmaco thérapeutique prescrit par le médecin est en conformité avec le protocole de traitement pour la réduction médicinale de paroxetine (Seroxat).

Le patient a reçu cette traitement pour la réduction des médicaments paroxetine (Seroxat) de la pharmacie Regenboog Apotheek et a réduit son médicament selon le calendrier.

Le coût des médicaments aux Pays-Bas (€ 77,-) sont répertoriés sur la facture.

Conforme l'arrêt Decker/Kohll (C-120/95, et C-158/06), le patient peut réclamer le remboursement des médicaments selon le taux applicable dans son pays natal.

(ES) Reembolso conforme a las sentencias C-120/95 y C-158/96 del Tribunal de Justicia de las Comunidades Europeas (sentencias Decker/Kohll)

El tratamiento farmacoterapéutico prescrito por el médico cumple con el protocolo de tratamiento para la supresión progresiva de paroxetine (Seroxat).

El paciente ha recogido en la farmacia Regenboog la medicación de supresión progresiva de paroxetine (Seroxat) y ha ido suprimiendo gradualmente la medicación según el esquema previsto.

El coste de la medicación en los Países Bajos (€ 77,-) consta en la factura.

Conforme a las sentencias Decker/Kohll (C-120/95 y C-158/06), el paciente tiene derecho al reembolso de la medicación según la tarifa vigente en su país de origen.

(DK) Erstatning ifølge C-120/95 og C-158/96 fra det Europæiske Fællesskabs domstol (Decker/Kohll-dommen)

Den af lægen ordinerede farmakologiske behandling er i overensstemmelse med behandlingsprotokollen for udfasning af paroxetine (Seroxat).

Patienten har modtaget nedtrapningsmedicin fra Regenboog Apotheek paroxetine (Seroxat) og har gradvist reduceret sin medicin ifølge skemaet.

Prisen på medicinen i Nederlandene (€ 77,-) er oplyst på fakturaen.

Ifølge Decker/Kohll-dommen (C-120/95 og C-158/06) er patienten berettiget til refusion af lægemidlet til den sats, der gælder i hjemlandet.

(PT) Reembolso conforme os Processos C-120/95 e C-158/96 do Tribunal de Justiça da Comunidade Europeia (Acórdão Decker/Kohll)

O tratamento farmacoterapêutico prescrito pelo médico está de acordo com o protocolo de tratamento para a redução de paroxetine (Seroxat).

O doente recebeu medicação de desabitação da Farmácia Regenboog e reduziu gradualmente a medicação, de acordo com o plano.

Os custos do medicamento na Holanda (€ 77,-) estão indicados na fatura.

De acordo com o Acórdão Decker/Kohll (Processos C-120/95 e C-158/06), o doente tem direito ao reembolso do medicamento à taxa aplicável no seu país de origem.

(NO) Refusjon i samsvar med den Europeiske Domstolens dom i sak C-120/95 (Decker/Kohll) og C-158/96.

Den farmakoterapeutiske behandlingen som er foreskrevet av praktiserende lege, samsvarer med behandlingsprotokollen for å redusere bruk av paroxetine (Seroxat).

Pasienten har fått paroxetine (Seroxat) for å redusere bruken av dette legemidlet og har gradvis redusert bruken i overensstemmelse med planen.

Kostnadene for medikamentene i Nederland (€ 77,-) er angitt på fakturaen.

I henhold til Decker/Kohll-dommen (C-120/95 og C-158/06) har pasienten rett til refusjon av kostnadene som gjelder for den medisinske behandling i pasientens hjemland.

(SE) Återbetaling enligt C-120/95 och C-158/96 från europadomstolen (Decker/Kohll-dom)

Den farmakoterapeutiska behandlingen, som föreskrevs av läkaren, motsvarar behandlingsprotokollet för att minska paroxetine (Seroxat).

Patienten har fått paroxetine (Seroxat)-medicinering för att minska användningen av detta läkemedel och har gradvis minskat användningen enligt schemat.

Kostnaderna för medicinen i Nederländerna (€ 77,-) står på fakturan.

I enlighet med Decker/Kohll- domen (C-120/95 sv C-158/06) har patienten rätt till återbetaling av priset för läkemedel.

(FI) Korvaus Euroopan yhteisön tuomioistuimen C-120/95 ja C-158/96 mukaan (tuomio Decker/Kohll)

Lääkäriin määräämä farmakoterapeuttinen hoito vastaa hoitomenetelmää tuotteen paroxetine (Seroxat) vähentämiseksi.

Potilas on saanut paroxetine (Seroxat)-lääkettä vähentääkseen tämän lääkkeen käyttöä ja vähentänyt käyttöä asteittain aikataulun mukaisesti.

Lääkityskustannukset Alankomaissa (€ 77,-) ilmoitetaan laskussa.

Decker/Kohll-tuomion (C-120/95 fi C-158/06) mukaan potilaalla on oikeus hänen kotimaassaan sovellettavaan korvaukseen lääkityksen hinnasta.

et i enlighet med det pris som tillämpas i hemlandet.

(PO) Zwrot kosztów jest zgodny z orzeczeniami Europejskiego Trybunału Sprawiedliwości w sprawach C-120/95 i C-158/96 (wyrokami w sprawach Decker/Kohll).

Leczenie farmakoterapeutyczne, które zostało przepisane przez lekarza, jest zgodne z protokołem leczenia mającego na celu zmniejszenie dawki leku paroxetine (Seroxat).

Pacjent otrzymywał lek paroxetine (Seroxat) w celu ograniczenia stosowania tego leku i stopniowo zmniejszał jego stosowanie zgodnie z harmonogramem.

Koszty leku w Holandii (€ 77,-) są podawane na fakturze.

Zgodnie z orzeczeniami w sprawach Decker/Kohll (C-120/95 i C-158/06), pacjent ma prawo do zwrotu kosztów leku według taryfy obowiązującej w jego kraju ojczystym.

(IT) Rimborso conforme alle sentenze della Corte di Giustizia Europea nei procedimenti C-120/95 e C-158/96 (sentenza Decker/Kohll)

Il trattamento farmacoterapeutico prescritto dal medico corrisponde al protocollo di trattamento per la riduzione di paroxetine (Seroxat).

Il paziente ha ricevuto il farmaco paroxetine (Seroxat) per ridurre l'uso di questo medicinale e ne ha gradualmente ridotto l'uso secondo il programma.

I costi del farmaco nei Paesi Bassi ammontanti a (€ 77,-) sono riportati in fattura.

In base alle sentenze emesse nei procedimenti Decker/Kohll (C-120/95 e C-158/06), il paziente ha diritto al rimborso dei costi sostenuti per l'acquisto del farmaco nel proprio paese di residenza.

(HU) Visszatérítés az Európai Bíróság C-120/95 és a C-158/96 ítélete szerint (Decker/Kohll ítélet)

A kezelőorvos előírta gyógyszeres kezelés megfelel a/az paroxetine (Seroxat) csökkentését irányzó protokollnak.

A páciens megkapta a/az paroxetine (Seroxat) gyógyszert ennek a gyógyszernek a fokozatos csökkentése érdekében, és az ütemterv szerint fokozatosan végezte a csökkentést.

Hollandiában a gyógyszer ára fel van tüntetve a számlán (€ 77,-).

A Decker/Kohll ítéletnek megfelelően (C-120/95 en C-158/06) a beteg jogosult a gyógyszer árának visszatérítésére a saját országában érvényes díjazás szerint.

(GR) Συμμόρφωση αποζημίωσης C-120/95 και C-158/96 από το δικαστήριο της Ευρωπαϊκής Εταιρείας (απόφαση Decker/Kohll)

Η φαρμακοθεραπευτική θεραπεία που ορίστηκε από τον ιατρό συμφωνεί με το πρωτόκολλο θεραπείας για τη μείωση paroxetine (Seroxat).

Ο ασθενής έχει λάβει φαρμακευτική αγωγή paroxetine (Seroxat) για μείωση της χρήσης του φαρμάκου αυτού και έχει μειώσει σταδιακά τη χρήση σύμφωνα με το χρονοδιάγραμμα.

Το κόστος της φαρμακευτικής αγωγής στις Κάτω Χώρες (€ 77,-) δημοσιεύεται στο τιμολόγιο.

Σύμφωνα με την απόφαση Decker/Kohll (C-120/95 και C-158/06) ο ασθενής δικαιούται επιστροφή εξόδων για την αγορά φαρμάκων σύμφωνα με το τιμολόγιο που ισχύει στη χώρα καταγωγής του.

(EE) Hüvitamine on kooskõlas Euroopa Kohtu otsustega C-120/95 ja C-158/96 (Decker/Kohlli otsused)

Praktiseeriva arsti välja kirjutatud farmakoterapeutiline ravi vastab vähendamise raviprotokollile paroxetine (Seroxat).

Patsient on saanud paroxetine (Seroxat) ravimeid selle ravimi kasutamise vähendamiseks ja järk-järgult vastavalt ajakavale vähendanud selle kasutamist.

Hollandis (€ 77,-) avaldatakse ravimikulud arvel.

Decker/Kohlli otsuse (C-120/95 ja C-158/06) kohaselt on patsiendil õigus saada ravikuuri eest tema koduriigis kehtivat hüvitist.

BG Възстановяването на разходите е съгласно решение по дело C-120/95 и C-158/96 на Съда на Европейския съюз (решение по дело Decker/Kohll)

Фармакотерапевтичното лечение, което е предписано от лекаря съответства на протокола за лечение с намаляване на дозата на paroxetine (Seroxat).

Пациентът е получил paroxetine (Seroxat) с цел намаляване на приема на този медикамент и постепенно е намалил приема съгласно схемата.

Стойността на медикамента в Холандия (€ 77,-) е посочена на фактурата.

Съгласно решението по дело Decker/Kohll (C-120/95 и C-158/06) пациентът има право на възстановяване на разходите за медикамента съгласно тарифата за лекарствата, приложима в държавата на пациента.

IS Samræmd endurgreiðsla C-120/95 og C-158/96 frá dómstól Evrópusamfélagsins (Decker/Kohll-dómur)

Lyfjameðferðin, sem ávísað var af lækni, er í samræmi við aðferðarlýsingu meðferðar til að draga úr paroxetine (Seroxat).

Sjúklingur hefurengið paroxetine (Seroxat) lyf til að draga úr notkun lyfsins og hefur smám saman dregið úr notkun samkvæmt áætlun.

Kostnaður við lyfið í Hollandi (€ 77,-) er birt á reikningnum.

Samkvæmt Decker/Kohll-dómnum (C-120/95 en C-158/06) á sjúklingurinn rétt á endurgreiðslu lyfsins samkvæmt samræmdri gjaldskrá lyfja sem við á í heimalandi hans.

RO Rambursare în conformitate cu C-120/95 și C-158/96 pronunțate de Curtea Europeană de Justiție (hotărârea Decker/Kohll)

Tratamentul farmacoterapeutic, care a fost prescris de medic, corespunde protocolului de tratament pentru scăderea dozei de paroxetine (Seroxat).

Pacientul a primit medicamentul paroxetine (Seroxat) pentru a reduce utilizarea acestui medicament și a redus treptat utilizarea conform programului.

Costurile medicamentelor din Olanda (€ 77,- Euro) sunt publicate pe factură.

Conform hotărârii Decker/Kohll (C-120/95 en C-158/06), pacientul are dreptul la rambursarea costului medicamentului aplicabil în țara de origine.

SK Náhrada je v súlade s C-120/95 a C-158/96 Európskeho súdneho dvora (Rozsudok Decker/Kohll)

Farmakoterapeutická liečba predpísaná lekárom je v súlade s protokolom o liečbe na zmiernenie paroxetine (Seroxat).

Pacient dostal liečivo paroxetine (Seroxat) na zníženie miery užívania tohto liečiva. Podľa harmonogramu mieru užívania znížil.

Ceny za liečivo v Holandsku (€ 77,- Euro) sú uvedené na faktúre.

Podľa rozsudku v prípade Decker/Kohll (C-120/95 a C-158/06) má pacient nárok na preplatenie liečiva v sadzbe, ktoré platí pre jeho domovskú krajinu.

CZ Úhrada výdajů odpovídající C-120/95 a C-158/96 ze Soudního dvoru Evropské společnosti (Decker/Kohll-judgment)

Farmakoterapeutická léčba, která byla předepsána praktickým lékařem, odpovídá protokolu léčby pro redukcii paroxetine (Seroxat).

Pacient obdržel lék paroxetine (Seroxat) ke snížení užívání tohoto léku a postupně užívání snížil podle plánu.

Náklady na léčbu v Holandsku (€ 77,- Euro) jsou zveřejněny na faktuře.

Podle rozsudku Decker/Kohll-judgment (C-120/95 en C-158/06) má pacient právo na náhradu léků odpovídající ceníku platnému v této domovské zemi.

HR Nadoknada troškova u skladu sa C-120/95 i C-158/96 sa Suda pravde Evropskog društva (presuda Decker/Kohll)

Farmakoterapijski tretman koji je propisao liječnik odgovara protokolu liječenja za smanjenje doze lijeka paroxetine (Seroxat).

Pacijent je primio lijek paroxetine (Seroxat) za smanjenje uporabe ovog lijeka i postupno je smanjivao upotrebu istog prema rasporedu.

Troškovi lijeka u Nizozemskoj (€ 77,- Euro) objavljeni su na računu.

Prema presudi Decker/Kohll (C-120/95 en C-158/06) pacijent ima pravo na nadoknadu tarife za lijekove koja je primjenjiva u njegovoj matičnoj zemlji.

SI Nadomestilo skladno z rzsodbama C-120/95 in C-158/96 Sodišča Evropske unije (sodba v primeru Decker/Kohll)

Farmakoterapevtsko zdravljenje, ki ga je predpisal zdravnik, je skladno s protokolom zdravljenja za zmanjšanje paroksetine (Seroxat).

Pacient je prejel zdravilo paroksetine (Seroxat) za zmanjšanje uporabe tega zdravila in je dejansko znatno zmanjšal uporabo skladno z razporedom.

Stroški zdravil na Nizozemskem (**€ 77,- evrov**) so navedeni na računu.

Skladno s sodbo Decker/Kohl (C-120/95 in C-158/06) je pacient upravičen do nadomestila za zdravila skladno z veljavno tarifo v svoji državi.

LV **Kompensāšana atbilst Eiropas Kopienas tiesas C-120/95 un C-158/96 noteikumiem (Decker/Kohl-judgment)**

Farmakoterapeitiskā terapija, kuru ordinējis ārsts, atbilst paroksetine (Seroxat) samazināšanas ārstēšanas protokolam.

Pacients ir saņēmis paroksetine (Seroxat) zāles, lai samazinātu šo zāļu devas lietošanu, un ir panācis pakāpenisku tās samazināšanu saskaņā ar grafiku.

Zāļu cenas Nīderlandē (**€ 77,- eiro**) ir norādītas rēķinā.

Saskaņā ar Dekera/Kola (Decker/Kohl) spriedumu (C-120/95 en C-158/06) pacientam ir tiesības uz zāļu kompensāciju atbilstoši zāļu cenai viņa mītnes zemē.

SUMMARY, 24 May 2018

Antidepressant tapering strips to help people come off medication more safely

Peter Groot and Jim van Os

Antidepressant users who have come to the end of treatment and wish to stop can suffer withdrawal symptoms, even when following their doctor's advice. In some cases, these withdrawal symptoms can be so severe that they prevent the user from stopping their medication. This can often be problematic for the patient and can result in unnecessary prescriptions, additional cost for health services and potential long-term health issues.

An observational study published in the journal *Psychosis* shows that antidepressant withdrawal symptoms can be minimised or avoided by using prepackaged medication in tapering strips. Tapering medication makes safe withdrawal possible by utilising pills of small dosages to make much smaller dosage reductions than can be achieved using standard registered dosages. This approach makes tapering easy for the antidepressant user and the doctor. The study shows that more patients are successful using tapering strips when compared to those who had tried to reduce using other methods and allows patients who have previously tried and failed to withdraw the opportunity to succeed in coming off their antidepressant medication.

What is tapering medication?

Tapering medication is prepackaged in tapering strips which make it easy for the patient to reduce without needing to follow complicated instructions. Furthermore the strips make it easy for doctors to prescribe personalised tapering schedules for their patients. A tapering strip consists of 28 sachets on a roll, with each sachet providing the medication for a single day. The daily dosage in each sachet is a little less or equal to that of the previous day. A doctor, psychiatrist or therapist can prescribe tapering schedules tailored to and in consultation with the patient in a shared decision-making approach. Using currently available standard registered dosages, gradual tapering is often difficult and, in some cases, impossible to achieve. Due to these limitations, current tapering guidelines are also difficult to follow for both the prescriber and the patient.

About the study

The study reviewed 1121 questionnaires completed by patients who wanted to withdraw from their antidepressant medications. The majority (75%, $n = 895$) wanted to taper off completely. Of these, 62% ($n = 692$) had previously made one or more tapering attempts. Almost everyone (97%) had suffered withdrawal symptoms during these earlier attempts, which were rated as very severe by 49% (7 on a scale of 1-7). The duration of antidepressant use varied from less than a year to more than 30 years (median: 2-5 years).

A comparison of withdrawal with or without tapering medication

All patients were asked about their tapering experience, using a scale from 1-7, where 1 = very good and 7 = very bad. In addition, they were asked if they had experienced withdrawal symptoms on a 1-7 scale, where 1 = not at all and 7 = very much. Patients were also asked to rate their previous attempts to withdraw, therefore making a comparison possible between tapering medication and other methods.

Study findings

The main results of the study are that, of the 895 people who used tapering strips to taper the antidepressant they used:

- 71% ($n = 636$) succeeded in a median time of 56 days (2 tapering strips).

- 21% (n = 192) did not complete the taper, 8% (n = 67) because they were still tapering, 4% (n = 39) due to withdrawal symptoms and 6% (n = 53) due to relapse (defined as the recurrence of complaints such as anxiety or depression).

The results of this study show that, for patients who had previously tried and failed to withdraw, the use of tapering medication in tapering strips could enable them to withdraw successfully.

Tapering medication enables shared decision making

For patients who wish to withdraw, attempting to taper using standard registered doses increases the risk of withdrawal symptoms, leading to an increased risk of withdrawal failure and unnecessary long-term antidepressant use. Allowing patients and doctors to work together to define personalised tapering schedules has been shown to reduce withdrawal symptoms and increase the chances of success. By doing this, doctors can focus on patient recovery and more easily distinguish between withdrawal symptoms and relapse, which is important in avoiding the unnecessary prescribing of antidepressant medications.

Applying tapering principles to other medications

The results of this study are relevant for people using other psychotropic medications such as antipsychotics or benzodiazepines. This is important for people with psychosis who will often be taking multiple medications. The use of tapering medication in tapering strips offers the opportunity to make safe, gradual dosage adjustments while minimising unpleasant withdrawal symptoms even in those patients who are taking multiple medications. The use of tapering strips enables a level of flexibility and personalisation that is simply not achievable using conventional standard registered dosages.

Information about taperingstrips can be found on the User Research Center of Maastricht University website www.taperingstrip.org

Dr. Peter C. Groot is a researcher and expert by experience at the User Research Center of Maastricht University and volunteer at the Cinderella Therapeutics Foundation. Prof. Jim van Os is Chairman of the Brain Division at the Utrecht University Medical Center.

Email: p.c.groot@maastrichtuniversity.nl, j.j.vanos-2@umcutrecht.nl.
